EGD (Endoscopy) Preparation Instructions

Date of Procedure:	Location:
Arrival Time:	Procedure Time:

- Arrange Ride Home
 - o You will not be able to drive yourself because of sedation. BART, taxi, limousines, and public transportation are not allowed.

When	Action	Examples
5 Days Before NO BLOOD THINNERS; NO ASPIRIN	Important: Medication change may be needed; read all of them. Confirm with your doctor if you need to stop any of your blood thinners No seeds such as tomato products or sesame seeds	 Plavix ,Aspirin, Ibuprofen, Naproxen, Advil, Motrin, Nuprin, Indomethacin, Celebrex, Aggrenox, etc-stop 5 days prior Coumadin – stop 5 days prior Lovenox, Heparin – do not take within 24 hours Pradaxa – stop 2 days prior (4 days if CKD) Xarelto/Eliquis/Savaysa – do not take within 24 hours (48 hours if CKD)
Day Before	 Eat and drink normally for breakfast and lunch light dinner Do not eat anything within 8 hours of your procedure. You can sip water for medicine 	 You can take your usual medications unless instructed otherwise by your physician with sips of water Modify diabetes medications if instructed by your doctor
Day of the Exam	Nothing by mouth 2 hours prior procedure.	 Wear loose comfortable clothing Leave valuables and jewelry at home Bring photo ID and insurance card Bring a complete list of medications and allergies
After endoscopy	Rest	No driving for the rest of the day or making major decisions.

If you need to cancel or reschedule, kindly give 5 business days to do so or there will be a \$100 fee; this will not be covered by your insurance company.

If you have any more questions, please call our office at (925)776-7600.